

Testing the Reliability of the Self-complementary Noncovalent Interactions: Supramolecular Implications and Supramolecular Design

Catalina Ruiz-Pérez, *Laboratorio de Rayos X y Materiales Moleculares, Departamento de Física Fundamental II, Universidad de La Laguna, La Laguna (Tenerife), Spain.* E-mail: caruiz@ull.es.

Noncovalent interactions play a special role in supramolecular chemistry, which has been defined by Lehn [1] as “chemistry beyond the molecule”. Noncovalently assisted synthetic procedures are used to assemble various types of supramolecular species. These syntheses rely on the stabilization provided by noncovalent interactions between recognition sites incorporated within precursors. As a recognition motif utilized to guide the synthesis, various types of noncovalent interactions can be used. These are, specifically, hydrogen bonds (H-bonds), stacking interactions, electrostatic interactions, hydrophobic interactions, charge-transfer interactions, and metal coordination [2]. Unconventional polymers composed of covalent and noncovalent bonds differ dramatically from standard, conventional polymers with just covalent bonds. They possess novel physical, optical, electrochemical, photochemical, biological, and catalytic properties. Targeted synthesis of macro- and supramolecular structures of various sizes, shapes, and functionality has now become possible. Supramolecular chemistry offers incredible applications in various fields such as medical chemistry (drug delivery systems), host-guest chemistry, catalysis and molecular electronics.

[1] Lehn J.-M., *Angew. Chem., Int. Ed. Engl.*, 1988, **27**, 89, *ibid.* 1990, **29**, 1304. [2] Lehn J.-M., Atwood J. L., Davies J. E. D., MacNicol D. D., Vögtle F., *Comprehensive Supramolecular Chemistry*, Eds. Pergamon, Oxford, 1996.

Keywords: supramolecular, noncovalent interactions, coordination compounds